

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 September 2003 (25.09.2003)

PCT

(10) International Publication Number
WO 03/078404 A1

(51) International Patent Classification⁷: **C07D 239/48**,
A61K 31/505, A61P 35/04

(21) International Application Number: PCT/EP03/02710

(22) International Filing Date: 14 March 2003 (14.03.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0206215.6 15 March 2002 (15.03.2002) GB

(71) Applicant (for all designated States except AT, US): **NOVARTIS AG** [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).

(71) Applicant (for AT only): **NOVARTIS PHARMA GMBH** [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BAENTELI, Rolf** [CH/CH]; Davidsbodenstrasse 64, CH-4056 Basel (CH). **ZENKE, Gerhard** [DE/DE]; Adolf-Gilattacker-Strasse 8, 79618 Rheinfelden (DE). **COOKE, Nigel, Graham** [GB/CH]; Tichelengraben 13, CH-4104 Oberwil (CH). **DUTHALER, Rudolf** [CH/CH]; Girenholdenweg 17, CH-4126 Bettingen (CH). **THOMA, Gebhard** [DE/DE]; Talweg 32, 79540 Lörrach (DE). **VON MATT, Anette** [DE/CH]; Fichtlirain 38, 4105 Biel-Benken (CH). **HONDA, Toshiyuki** [JP/JP]; 20-17, Inarimae, Tsukuba-shi, Ibaraki Pref. 305-0061 (JP). **MATSUURA,**

Naoko [JP/JP]; 2-4-1-507, Amakubo, Tsukuba-shi, Ibaraki Pref. 305-0005 (JP). **NONOMURA, Kazuhiko** [JP/JP]; 1-13-9-603, Sakura, Tsukuba-shi, Ibaraki-Pref. 305-0003 (JP). **OHMORI, Osamu** [JP/JP]; 2-4-6-406, Sengen, Tsukuba-shi, Ibaraki Pref. 305-0047 (JP). **UMEMURA, Ichiro** [JP/JP]; 2-3-7-406, Ninomiya, Tsukuba-shi, Ibaraki Pref. 305-0051 (JP). **HINTERDING, Klaus** [DE/DE]; Lörracherstrasse 92, 79595 Rümmingen (DE). **PAPAGEORGIOU, Christos** [FR/FR]; 25b, rue de Castelnau, F-68400 Riedisheim (FR).

(74) Agent: **GROS, Florent**; Novartis AG, Corporate Intellectual Property, CH-4002 Basel (CH).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LT, LU, LV, MA, MD, MK, MN, MX, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SE, SG, SK, TJ, TM, TN, TR, TT, UA, US, UZ, VC, VN, YU, ZA, ZW.

(84) Designated States (regional): Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR).


Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 03/078404 A1

(54) Title: PYRIMIDINE DERIVATIVES




(I)

(57) Abstract: There are provided compounds of formula (I), wherein X, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸ and R⁹ are as indicated in claim 1, useful in disorders where ZAP-70 and/or Syk inhibition plays a role or caused by a malfunction of signal cascades connected with FAK.

2,4-DIAMINOPYRIMIDINE DERIVATIVES

The present invention relates to pyrimidine derivatives, to processes for their production, their use as pharmaceuticals and to pharmaceutical compositions comprising them.

More particularly the present invention provides in a first aspect, a compound of formula I



wherein

X is =CR⁰- or =N-;

each of R⁰, R¹, R², R³ and R⁴ independently is hydrogen; hydroxy; C₁-C₈alkyl; C₂-C₈alkenyl;

C₃-C₈cycloalkyl; C₃-C₈cycloalkyl-C₁-C₈alkyl; hydroxyC₁-C₈alkyl; C₁-C₈alkoxyC₁-C₈alkyl; hydroxyC₁-C₈alkoxyC₁-C₈alkyl; arylC₁-C₈alkyl which optionally may be substituted on the ring by hydroxy, C₁-C₈alkoxy, carboxy or C₁-C₈alkoxycarbonyl;

or R³ and R⁴ form together with the nitrogen and carbon atoms to which they are attached a 5 to 10 membered heterocyclic ring and comprising additionally 1, 2 or 3 heteroatoms selected from N, O and S;

or each of R¹, R² and R³, independently, is halogen; halo-C₁-C₈alkyl; C₁-C₈alkoxy; halo-C₁-C₈alkoxy; hydroxyC₁-C₈alkoxy; C₁-C₈alkoxyC₁-C₈alkoxy; aryl; arylC₁-C₈alkoxy; heteroaryl; heteroaryl-C₁-C₄alkyl; 5 to 10 membered heterocyclic ring; nitro; carboxy; C₂-C₈alkoxycarbonyl; C₂-C₈alkylcarbonyl; -N(C₁-C₈alkyl)C(O)C₁-C₈alkyl; -N(R¹⁰)R¹¹; -CON(R¹⁰)R¹¹; -SO₂N(R¹⁰)R¹¹; or -C₁-C₄-alkylene-SO₂N(R¹⁰)R¹¹; wherein each of R¹⁰ and R¹¹ independently is hydrogen; hydroxy; C₁-C₈alkyl; C₂-C₈alkenyl; C₃-C₈cycloalkyl; C₃-C₈cycloalkyl-C₁-C₈alkyl; C₁-C₈alkoxyC₁-C₈alkyl; hydroxyC₁-C₈alkoxyC₁-C₈alkyl; hydroxyC₁-C₈alkyl; (C₁-C₈alkyl)-carbonyl; arylC₁-C₈alkyl which optionally may be substituted on the ring by hydroxy, C₁-C₈alkoxy, carboxy or C₂-C₈alkoxycarbonyl; or 5 to 10 membered heterocyclic ring;

or R¹ and R² form together with the C-atoms to which they are attached aryl or a 5 to 10 membered heteroaryl residue comprising one or two heteroatoms selected from N, O and S; or

each of R⁵ and R⁶ independently is hydrogen; halogen; cyano; C₁-C₈alkyl; halo-C₁-C₈alkyl;

C₂-C₈alkenyl; C₂-C₈alkynyl; C₃-C₈cycloalkyl; C₃-C₈cycloalkylC₁-C₈alkyl; C₅-C₁₀arylC₁-C₈alkyl; each of R⁷, R⁸ and R⁹ is independently hydrogen; hydroxy; C₁-C₈alkyl; C₂-C₈alkenyl; halo-C₁-C₈alkyl; C₁-C₈alkoxy; C₃-C₈cycloalkyl; C₃-C₈cycloalkylC₁-C₈alkyl; arylC₁-C₈alkyl; -Y-R¹² wherein Y is a direct bond or O and R¹² is a substituted or unsubstituted 5, 6 or 7 membered heterocyclic ring comprising 1, 2 or 3 heteroatoms selected from N, O and S; carboxy; (C₁-C₈alkoxy)-carbonyl; -N(C₁₋₈alkyl)-CO-NR¹⁰R¹¹; -CONR¹⁰R¹¹; -N(R¹⁰)(R¹¹); -SO₂N(R¹⁰)R¹¹; or R⁷ and R⁸ or R⁸ and R⁹, respectively form together with the carbon atoms to which they are attached, a 5 or 6 membered heteroaryl comprising 1, 2 or 3 heteroatoms selected from N, O and S; or a 5 or 6 membered carbocyclic ring.

in free form or salt form.

Any aryl may be phenyl, naphthyl or 1,2,3,4-tetrahydronaphthyl, preferably phenyl. Heteroaryl is an aromatic heterocyclic ring, e.g. a 5 or 6 membered aromatic heterocyclic ring, optionally condensed to 1 or 2 benzene rings and/or to a further heterocyclic ring.

Any heterocyclic ring may be saturated or unsaturated and optionally condensed to 1 or 2 benzene rings and/or to a further heterocyclic ring.

Examples of heterocyclic rings or heteroaryl include e.g. morpholinyl, piperazinyl, piperidyl, pyrrolidinyl, pyridyl, purinyl, pyrimidinyl, N-methyl-aza-cycloheptan-4-yl, indolyl, quinolinyl, isoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, benzothiazolyl, thiazolyl, imidazolyl, benzimidazolyl, benzoxadiazolyl, benzotriazolyl, indanyl, oxadiazolyl, pyrazolyl, triazolyl, and tetrazolyl. Preferred heterocyclic rings or heteroaryl are morpholinyl, piperazinyl, piperidyl, pyrrolidinyl, pyridyl, N-methyl-aza-cycloheptan-4-yl, thiazolyl, imidazolyl and tetrazolyl.

When R⁷ and R⁸ or R⁸ and R⁹ form together with the carbon atoms to which they are attached a 5 or 6 membered carbocyclic ring, this may preferably be cyclopentyl or cyclohexyl.

Halo-alkyl is alkyl wherein one or more H are replaced by halogen, e.g. CF₃.

Any alkyl or alkyl moiety may be linear or branched. C₁₋₈alkyl is preferably C₁₋₄alkyl. C₁₋₈alkoxy is preferably C₁₋₄alkoxy. Any alkyl, alkoxy, alkenyl, cycloalkyl, heterocyclic ring, aryl or heteroaryl may be, unless otherwise stated, unsubstituted or substituted by one or more substituents selected from halogen; OH; C₁-C₈alkyl; C₁-C₈alkoxy; nitro; cyano; COOH; carbamoyl;

$C(NH_2)=NOH$; $-N(R^{10})R^{11}$; C_3-C_6 cycloalkyl; 3 to 7 membered heterocyclic ring; phenyl; phenyl- C_{1-4} alkyl; 5 or 6 membered heteroaryl. When alkyl, alkoxy or alkenyl is substituted, the substituent is preferably on the terminal C atom. When the heterocyclic ring or heteroaryl is substituted, e.g. as disclosed above, this may be on one or more ring carbon atoms and/or ring nitrogen atom when present. Examples of a substituent on a ring nitrogen atom are e.g. C_{1-8} alkyl, carbamoyl, $-C(NH_2)=NOH$, $-NR^{10}R^{11}$, C_3-6 cycloalkyl or phenyl- C_{1-4} alkyl, preferably C_{1-8} alkyl, C_3-6 cycloalkyl or phenyl- C_{1-4} alkyl.

Preferably substituted alkyl or alkoxy as R_7 is alkyl or alkoxy substituted on the terminal C atom by OH, C_{1-4} alkoxy or a heterocyclic ring. When R^{10} or R^{11} is a 5 to 10 membered heterocyclic ring, it may be e.g. thiazolyl.

Halogen may be F, Cl, Br, or I.

Preferably at most one of R^1 , R^2 or R^3 is $CONR^{10}R^{11}$ or $SO_2NR^{10}R^{11}$, more preferably $SO_2NR^{10}R^{11}$.

The compounds of the invention may exist in free form or in salt form, e.g. addition salts with e.g. organic or inorganic acids, for example trifluoroacetic acid or hydrochloride acid, or salts obtainable when they comprise a carboxy group, e.g. with a base, for example alkali salts such as sodium, potassium, or substituted or unsubstituted ammonium salts.


In formula I the following significances are preferred independently, collectively or in any combination or sub-combination:

- (a) X is $=CR^0$;
- (b) R^0 is hydrogen; halogen, e.g. Cl; C_1-C_4 alkyl, e.g. methyl or ethyl; C_{1-4} alkoxy, e.g. methoxy; preferably hydrogen;
- (c) R^1 is hydrogen; halogen, e.g. Cl or F; OH; C_1-C_8 alkyl, e.g. methyl or ethyl; substituted C_{1-8} alkyl, e.g. terminally OH substituted C_{1-8} alkyl; $-SO_2N(R^{10})R^{11}$; $-N(C_{1-4}alkyl)C(O)C_{1-4}alkyl$; a 5 or 6 membered heterocyclic ring optionally substituted on a ring N atom (when possible); C_1-C_8 alkoxy, e.g. methoxy; aryl, e.g. phenyl; or form together with R^2 and the C-atoms to which R^1 and R^2 are attached 5 to 10 membered aryl or heteroaryl, the latter comprising 1 or 2 nitrogen atoms;

- (d) R^2 is hydrogen; hydroxy; C_1 - C_8 alkyl, e.g. methyl or ethyl; substituted C_{1-8} alkyl, e.g. terminally OH- or C_{1-4} -alkoxy substituted C_{1-8} alkyl; C_{1-8} alkoxy; C_1 - C_4 alkoxy C_{1-8} alkoxy; $-\text{CON}(R^{10})R^{11}$; $-\text{SO}_2\text{N}(R^{10})R^{11}$; or forms together with R^1 and the C-atoms to which R^1 and R^2 are attached a 5 to 10 membered aryl or heteroaryl, the latter comprising 1 or 2 nitrogen atoms;
- (e) R^3 is hydrogen; halogen, e.g. Cl, Br; hydroxy; C_1 - C_8 alkyl, e.g. methyl or ethyl; substituted C_{1-8} alkyl, e.g. terminally OH substituted C_{1-8} alkyl; carboxy; $\text{CONR}^{10}R^{11}$; $-\text{SO}_2\text{N}(R^{10})R^{11}$; a 5 or 6 membered heterocyclic ring optionally substituted on a ring nitrogen atom (when possible); or forms together with R^4 and the N and C atoms to which R^3 and R^4 are attached a 6 membered heterocyclic ring;
- (f) R^4 is hydrogen; or forms together with R^3 and the N and C atoms to which R^3 and R^4 are attached a 6 membered heterocyclic ring; preferably hydrogen;
- (g) R^5 is hydrogen; halogen; C_{1-4} alkyl; or CF_3 ;
- (h) R^6 is hydrogen;
- (i) R^7 is hydrogen; hydroxy; C_{1-4} alkyl; substituted C_{1-4} alkyl, e.g. terminally OH substituted C_{1-4} alkyl; C_{1-8} alkoxy; substituted C_{1-8} alkoxy, e.g. terminally substituted by OH, C_{1-4} alkoxy or a heterocyclic ring; $\text{NR}^{10}R^{11}$; $-\text{SO}_2\text{N}(R^{10})R^{11}$; $-\text{Y}-R^{12}$; CF_3 ; or R^7 forms together with R^8 and the C-atoms to which R^7 and R^8 are attached a 5 membered heteroaryl residue, e.g. bridged by $-\text{NH}-\text{CH}=\text{CH}-$, $-\text{CH}=\text{CH}-\text{NH}-$, $-\text{NH}-\text{N}=\text{CH}-$, $-\text{CH}=\text{N}-\text{NH}-$, $-\text{NH}-\text{N}=\text{N}-$ or $-\text{N}=\text{N}-\text{NH}-$;
- (k) R^8 is hydrogen; hydroxy; C_{1-4} alkoxy; carboxy; a 5 or 6 membered heterocyclic ring optionally substituted on a ring C or N atom; $\text{N}(C_{1-4}\text{alkyl})-\text{CO}-\text{NR}^{10}R^{11}$; or forms with R^7 or R^9 and the C-atoms to which R^7 and R^8 or R^8 and R^9 , respectively, are attached a 5 membered heteroaryl residue, e.g. bridged by $-\text{NH}-\text{CH}=\text{CH}-$, $-\text{CH}=\text{CH}-\text{NH}-$, $-\text{NH}-\text{N}=\text{CH}-$, $-\text{CH}=\text{N}-\text{NH}-$, $-\text{NH}-\text{N}=\text{N}-$ or $-\text{N}=\text{N}-\text{NH}-$;
- (l) R^9 is hydrogen; C_{1-4} alkoxy; $\text{NR}^{10}R^{11}$; or forms with R^8 and the C atoms to which R^8 and R^9 are attached a 5 membered heteroaryl, e.g. bridged by $-\text{NH}-\text{CH}=\text{CH}-$, $-\text{CH}=\text{CH}-\text{NH}-$, $-\text{NH}-\text{N}=\text{CH}-$, $-\text{CH}=\text{N}-\text{NH}-$, $-\text{NH}-\text{N}=\text{N}-$ or $-\text{N}=\text{N}-\text{NH}-$;
- (m) one of R^{10} and R^{11} , independently, is hydrogen or C_{1-4} alkyl and the other is hydrogen; OH; C_{1-8} alkyl, substituted C_{1-8} alkyl, e.g. terminally substituted by OH, C_{3-6} cycloalkyl or a heterocyclic ring; C_{2-8} alkenyl; C_{3-8} cycloalkyl; hydroxy C_{1-8} alkoxy C_{1-8} alkyl; or a 5 membered heterocyclic ring.

R^3 is preferably $\text{SO}_2\text{NR}^{10}R^{11}$.

The present invention also provides a process for the production of a compound of formula I, comprising reacting a compound of formula II



wherein R¹, R², R³, R⁴, R⁵, R⁶ and X are as defined above, and Y is a leaving group, preferably halogen such as bromide, iodine, or in particular chloride;

with a compound of formula III



wherein R⁷, R⁸ and R⁹ are as defined above;


and recovering the resulting compound of formula I in free or in form of a salt, and, where required, converting the compound of formula I obtained in free form into the desired salt form, or vice versa.

The process may be performed according to methods known in the art, e.g. as described in examples 1 to 4.

The compound of formula II used as starting materials may be obtained by reacting a compound of formula IV



with a compound of formula V




(V)

wherein R¹, R², R³, R⁴, R⁵, R⁶, Y and X are as defined above.

The compounds of formula IV and V are known or may be produced in accordance with known procedures.

The following examples illustrate the invention without any limitation.

The following abbreviations are employed: APC = allophycocyanine, BINAP = 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl, cDNA = complementary DNA, DCM = dichloromethane, DIAD = diisopropyl azodicarboxylate, DMAP = 4-dimethylaminopyridine, DMF = dimethylformamide, DMSO = dimethylsulfoxide, DMF = dimethylformamide; Pmc = 2,2,5,7,8-pentamethylchroman; tBu = *tert*-butyl; DIPCDI = N,N'-diisopropylcarbodiimid; DTT = 1,4-dithio-D,L-treitol, DNA = deoxyribonucleic acid, EDTA = ethylenediaminetetra-acetic acid, Lck = lymphoid T-cell protein tyrosine kinase, LAT-11 = linker for activation of T cell, RT = room temperature; RT-PCR = reverse transcription polymerase chain reaction, MS = molecular ion (e.g. M+H¹⁺) determined by electrospray mass spectroscopy; Eu = europium; ZAP-70 = zeta chain-associated protein of 70 kD; Syk = p72syk protein tyrosine kinase; SA = streptavidin.

Example 1: 2-[2-(1H-Indazol-6-ylamino)-pyrimidin-4-ylamino]-benzenesulfonamid


(a) *2-(2-Chloro-pyrimidin-4-ylamino)-benzenesulfonamide*: To a suspension of 8.52 g (49.47 mmol) 2-aminobenzenesulfonamide in 200 ml isopropanol is added 22.1 g (148.42 mmol, 3 equivalent) 2,4-dichloropyrimidine and 20 ml 10 M hydrochloric acid (200 mmol, 4 equivalent). The suspension is stirred at 60°C for 2 h 15 min. The reaction mixture is diluted with 2 l ethyl acetate and 500 ml water is added. The pH is adjusted to 8-9 by addition of sodium bicarbonate. The layers are separated and the aqueous layer is reextracted with 500 ml ethyl acetate. The organic layers are dried with sodium sulfate, filtered and evaporated to a volume of 300 ml. A crystalline precipitate is formed and removed by filtration (side product). The filtrate is evaporated to 100 ml whereupon the product crystallizes to give 2-(2-chloro-pyrimidin-4-ylamino)-benzenesulfonamide (97% purity by HPLC). The mother liquor of this crystallisation is further purified by column chromatography and crystallisation to give further 2-(2-chloro-pyrimidin-4-ylamino)-benzenesulfonamide.

(b) *2-[2-(1H-Indazol-6-ylamino)-pyrimidin-4-ylamino]-benzenesulfonamide*: To a suspension of 7.25 g (25.46 mmol) 2-(2-Chloro-pyrimidin-4-ylamino)-benzenesulfonamide and 4.07 g (30.55 mmol, 1.2 equivalent) 6-aminoindazole in 400 ml isopropanol is added 13 ml conc. HCl* (130 mmol, 5 equivalent). The suspension is refluxed for 4 h 30 min. The reaction mixture is diluted with 1.5 l ethyl acetate and 1 l water is added. The pH is adjusted to 8-9 by addition of sodium bicarbonate. The layers are separated and the aqueous layer is re-extracted with 500 ml ethyl acetate. The organic layers are dried with sodium sulfate, filtered and evaporated to a volume of 300 ml. A crystalline precipitate (1.01 g) is formed and removed by filtration (side product). The filtrate is purified by chromatography on 200 g silica gel eluting with ethyl acetate/methanol 95/5 v/v. Upon evaporation crystals are formed which are filtered to give the title compound.

¹H NMR (400 MHz, DMSO-d₆): δ 9.42 (s, 1H), 8.34 (d, 1H), 8.28 (d, 1H), 8.27 (s, 1H), 7.93 (s, 1H), 7.88 (d, 1H), 7.62 (m, 2H), 7.32 (d, 1H), 7.24 (t, 1H), 6.40 (d, 1H).

MS *m/z* (%): 382 (M+H, 100);


Example 2: 2-[2-(3,4,5-Trimethoxy-phenylamino)-pyrimidin-4-ylamino]-benzenesulfonamide



The title compound is prepared from 2-(2-chloro-pyrimidin-4-ylamino)-benzenesulfonamide as described in Example 1 using 3,4,5-Trimethoxy-phenylamine instead of 6-aminoindazole in step (b).

^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 9.18 (s, 1H), 8.22 (d, 1H), 8.17 (d, 1H), 7.89 (d, 1H), 7.55 (t, 1H), 7.25 (t, 1H), 7.14 (s, 2H), 6.40 (d, 1H), 3.69 (s, 6H), 3.62 (s, 3H). MS m/z (%): 432 ($\text{M}+\text{H}$, 100);

Example 3: 2-methyl-6-[2-(3,4,5-Trimethoxy-phenylamino)-pyrimidin-4-ylamino]-benzenesulfonamide



The title compound is prepared as described in Example 1 with the difference that in step (a) 2-amino-6-methyl-benzenesulfonamide is used instead of 2-aminobenzenesulfonamide.

2-Amino-6-methyl-benzenesulfonamide may be prepared as described by Girard, Y et al.; J. J. Chem. Soc. Perkin Trans. I 1979, 4, 1043-1047: Under an atmosphere of nitrogen *m*-toluidin (32.1 g, 32.5 ml, 0.30 mmol) is added dropwise to a solution of chlorosulfonyl isocyanate (51.3 ml, 83.6 g, 0.59 mmol) in nitroethane (400 ml) at $-55 - 49^\circ\text{C}$. The cold bath is removed and the mixture allowed to warm to -8°C , whereupon aluminium chloride (51 g, 0.38 mmol) is added. Heating the mixture to 100°C for 20 min forms a clear brown solution, which is cooled to RT and poured on ice. After filtration, washing with ice water and diethyl ether the precipitate is collected and dissolved in dioxane (300 ml). Water (1000 ml) and conc. HCl (1500 ml) are added to form a suspension, which is heated to 120°C for 18h. After cooling to RT the clear brown solution is washed with diethyl ether/hexane (1400 ml, 1/1 v/v) and adjusted to pH = 8 by addition of

sodium carbonate. Extraction using ethyl acetate (2 x 1000 ml), washing of the organic phase with water (500 ml) and brine (500 ml), drying (magnesium sulfate) and concentration yields a brown solid, which is purified by chromatography on silica using methylene chloride/ethanol (100/1 v/v) to yield the desired product as a white solid.


Melting point: 72-75°C (Propan-2-ol);

¹H NMR (400 MHz, DMSO-d₆): δ 2.64 (s, 3H, Me), 3.63 (s, 3H, OMe), 3.68 (s, 6H, OMe), 6.31 (d, J = 5Hz, 1H, pyrimidine CH), 7.07 (d, J = 8Hz, 1H, arom. CH), 7.15 (s, 2H, arom. CH), 7.40 (t, J = 8Hz, 1H, arom. CH), 7.65 (s, 2H, SO₂NH₂), 8.04 (d, J = 8Hz, 1H, arom. CH), 8.12 (d, J = 5Hz, 1H, pyrimidine CH), 9.14 (s, 1H, NH), 9.40 (s, 1H, NH).

MS (ES⁺) *m/z* 446 (MH⁺), 468 (MNa⁺)

MS (ES⁻): 444 (M-H)⁻

Example 4: 2-Methoxy-6-[2-(3,4,5-trimethoxy-phenylamino)-pyrimidin-4-ylamino]-benzenesulfonamide




The title compound is prepared as described in Example 1 with the difference that in step (a) 2-amino-6-methoxy-benzenesulfonamide is used instead of 2-Amino-6-methyl-benzenesulfonamide.

2-Amino-6-methoxy-benzenesulfonamide may be prepared from 12.3 g of meta-anisidine following an analogous procedure as described in Example 1a. NMR (400 MHz, DMSO-d₆): δ 3.62 (s, 3H, OMe), 3.69 (s, 6H, OMe), 3.91 (s, 3H, OMe), 6.31 (d, J = 5Hz, 1H, pyrimidine CH), 6.86 (d, J = 8Hz, 1H, arom. CH), 7.12 (s, 2H, arom. CH), 7.43 (t, J = 8Hz, 1H, arom. CH), 8.01 (d, J = 8Hz, 1H, arom. CH), 8.11 (d, J = 5Hz, 1H, pyrimidine CH), 9.18 (s, 1H, NH), 9.79 (br, 1H, NH).

MS (ES⁺): 462.2 (MH⁺), 484.2 (MNa⁺)

MS (ES⁻): 460.3 (M-H)⁻

The compounds of formula X₁




wherein R³, R⁷ and R⁸ are as defined in Table 1, may be prepared by following the procedure of Example 1 but using the appropriate starting materials.

TABLE 1

Example	R ³	R ⁷	R ⁸	MS Data		
				*ES+	*ES-	*EI
5	-OH	-O-(1-methyl)-azacyclohept-4-yl	-H	406	404	
6	-SO ₂ NH ₂	-O-(1-methyl)-azacyclohept-4-yl	-H	469.3		
7	-SO ₂ NH ₂	-O-2-(1-methyl-azacyclopent-2-yl)-ethyl	-H	469.3		
8	-OH	-O-2-(1-piperidyl)-ethyl	-OCH ₃	436.3	434.4	
9	-OH	-O-2-(1-methyl-azacyclopent-2-yl)-ethyl	-H	406	404	
10	-SO ₂ NH ₂	-O-CH ₂ CH ₂ CH ₂ -1-imidazolyl	-OCH ₃	496	494	
11	-SO ₂ NH ₂	-O-2-(1-piperidyl)-ethyl	-OCH ₃	499.2	497.3	
12	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -1-methyl-imidazol-1-yl	-H	466	464	
13	-OH	-O-2-[1-(1,2,4-triazolyl)]-ethyl	-H	390	388	
14	-OH	-O-2-hydroxyethyl	-OCH ₃	369.4	367.3	
15	-SO ₂ NH ₂	-O-2-hydroxyethyl	-OCH ₃			431
16	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -1-imidazolyl	-OCH ₃			
17	-SO ₂ NH ₂	-O-2-[1-(1,2,4-triazolyl)]-ethyl	-H			452
18	-SO ₂ NH ₂	-NH-N=N-			381	
19	-SO ₂ NHCH ₃	-O-CH ₂ CH ₂ -1-imidazolyl	-OCH ₃	496	494	
20	-SO ₂ NH ₂	-O-2-(1-piperidyl)-ethyl	-H	469	467	
21	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -1-imidazolyl	-H	452	450	
22	-OH	-O-2-(1-piperidyl)-ethyl	-H	406		
23	-COOH	-4-morpholino	-H			
24	-OH	-O-CH ₂ CH ₂ CH ₂ -1-imidazolyl	-OCH ₃	433	431	
25	-SO ₂ NHCH ₃	-CH=N-NH-		396	394	
26	-SO ₂ NH ₂	-O-2-(4-morpholino)ethyl	-H	471	469	
27	-SO ₂ NH ₂	-OCH ₃	-OCH ₃	402	400	
28	-OH	-O-2-(4-morpholino)ethyl	-H	408	406	
29	-SO ₂ NH ₂	-CH=N-NH-				381
30	-SO ₂ NHCH ₃	-O-CH ₂ CH ₂ -1-imidazolyl	-H			
31	-COOH	amino	-H	322		
32	-SO ₂ NH ₂	-O-CH ₂ CH ₂ CH ₂ -1-imidazolyl	-H	466.2	464.3	

33	-COOH	-N(CH ₃) ₂	-H			
34	-5-(1,2,3,4-tetrazolyl)	-NH-C(O)CH ₃	-H	388	386	
35	-SO ₂ NHCH ₃	-NH-N=CH-				
36	-COOH	-OH	-H			
37	-COOH	-H	-4-piperidyl			
38	-COOH	-CH ₂ -OH	-H			
39	-OH	-O-CH ₂ CH ₂ -1-imidazolyl	-OCH ₃			
40	-SO ₂ NH-CH ₂ CH ₂ -OH	-O-CH ₂ CH ₂ -1-imidazolyl	-H	496	494	
41	-C(O)NH ₂	amino	-H	321		
42	-SO ₂ NH ₂	-CH=CH-NH-		381		
43	-5-(1,2,3,4-tetrazolyl)	-NHCH ₂ -3-pyridyl	-H		435	
44	-SO ₂ NH ₂	-NH-CH=CH-			379	
45	-COOH	-H	-4-morpholino			
46	-COOH	-H	-1-(4-amino)-piperidyl			
47	-SO ₂ NH ₂	-OCH ₃	-H	372	370	
48	-SO ₂ N(CH ₃) ₂	-O-CH ₂ CH ₂ -1-imidazolyl	-H	480	478	

The compounds of formula X₂




wherein R³ and R⁸ are as defined in Table 2, may be prepared by following the procedure of Example 1 but using the appropriate starting materials.

TABLE 2

Example	R ³	R ⁸	MS Data	
			*ES+	*ES-
49	-COOH	-OCH ₃	397	395
50	-SO ₂ NH ₂	-OH		
51	-SO ₂ NHCH ₃	-OCH ₃		
52	-5-(1,2,3,4-tetrazolyl)	-OCH ₃	421	
53	-SO ₂ NH-cyclopropyl	-OCH ₃	472.2	470.3
54	-C(O)NHOH	-OCH ₃	412	410
55	-SO ₂ NH-CH ₂ CH ₂ -OH	-OCH ₃	476	474
56	-SO ₂ N(CH ₃) ₂	-OCH ₃	460.3	458.3

57	-OH	-OCH ₃	369	367
58	-SO ₂ NH-CH ₂ CH ₂ CH ₃	-OCH ₃	474	472
59	-CH ₂ OH	-OCH ₃		
60	-SO ₂ NH ₂	-H	402	

The compounds of formula X₃




wherein R¹, R⁷, R⁸ and R⁹ are as defined in Table 3, may be prepared by following the procedure of Example 1 but using the appropriate starting materials.

TABLE 3

Example	R ¹	R ⁷	R ⁸	R ⁹	MS Data	
					*ES+	*ES-
61	-SO ₂ NH-CH ₂ CH ₂ -O-CH ₂ CH ₂ -OH	-H	-N(CH ₃)-C(O)CH ₃	-H		
62	-SO ₂ NH ₂	-OCH ₃	-OCH ₃	-OCH ₃		
63	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -1-imidazolyl	-OCH ₃	-H		
64	-SO ₂ NH-CH ₂ CH ₂ -O-CH ₂ CH ₂ -OH	-OCH ₃	-OCH ₃	-OCH ₃	520	518
65	-N(CH ₃) C(O)CH ₃	-OCH ₃	-OCH ₃	-OCH ₃	424	422
66	-CH ₂ CH ₂ -OH	-SO ₂ NH-CH ₂ CH ₂ CH ₂ CH ₃	-H	-H		
67	-SO ₂ NH ₂	-OCH ₃	-H	-OCH ₃		
68	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -1-imidazolyl	-H	-H		
69	-CH ₂ CH ₂ -OH	-O-CH ₂ CH ₂ -1-imidazolyl	-H	-H		
70	-CH ₂ CH ₂ -OH	-OCH ₃	-H	-OCH ₃		
71	-SO ₂ NH ₂	-OH	-H	-H		
72	-O-CH ₂ CH ₂ -OH	-O-CH ₂ CH ₂ -1-imidazolyl	-H	-H		
73	-SO ₂ NH-2-thiazolyl	-OCH ₃	-OCH ₃	-OCH ₃	515	513

The compounds of formula X₄




wherein R², R⁵, R⁷, R⁸ and R⁹ are as defined in Table 4, may be prepared by following the procedure of Example 1 but using the appropriate starting materials.

TABLE 4

Example	R ²	R ⁵	R ⁷	R ⁸	R ⁹	MS Data	
						*ES+	*ES-
74	-SO ₂ NH-2-propenyl	-H	-OCH ₃	-OCH ₃	-OCH ₃	472	470
75	-SO ₂ NH ₂	-H	-OCH ₃	-OCH ₃	-OCH ₃		
76	-OH	-H	-O-(1-methyl)-azacyclohept-4-yl	-H	-H	406.3	404.3
77	-OH	-H	-O-CH ₂ CH ₂ -OH	-OCH ₃	-H	369	367
78	-SO ₂ NH ₂	-Br	-OCH ₃	-OCH ₃	-OCH ₃	510.1/ 512.1	508.1/ 510.2
79	-SO ₂ NH ₂	-H	-CH=N-NH-		-H	382	
80	-SO ₂ NH ₂	-CH ₃	-OCH ₃	-OCH ₃	-OCH ₃	446	444
81	-SO ₂ NH ₂	-H	-O-CH ₂ CH ₂ -1-imidazolyl	-OCH ₃	-H	482	480
82	-OH	-H	-O-CH ₂ CH ₂ -1-piperidyl	-OCH ₃	-H	436.3	434.3
83	-OH	-H	-O-CH ₂ CH ₂ -1-imidazolyl	-OCH ₃	-H	419	417
84	-SO ₂ NH ₂	-H	-O-CH ₂ CH ₂ -1-imidazolyl	-H	-H	452	450
85	-CH ₃	-C≡N	-OCH ₃	-OCH ₃	-OCH ₃	392	
86	-SO ₂ NH ₂	-H	-NH-N=CH-		-H	382	
87	-OH	-H	-OCH ₃	-OCH ₃	-OCH ₃	369	367
88	-SO ₂ NHCH ₃	-CH ₃	-OCH ₃	-OCH ₃	-OCH ₃	460	458
89	-OH	-H	-OH	-COOH	-OCH ₃		
90	-OH	-H	-O-CH ₂ CH ₂ -1-piperidyl	-H	-H	406	404
91	-SO ₂ NH-2-propenyl	-H	-O-CH ₂ CH ₂ -1-imidazolyl	-H	-H	492.3	490.3
92	-SO ₂ NH ₂	-Br	-O-CH ₂ CH ₂ -1-(1-methyl)-imidazolyl	-H	-H	544.1/ 546	542/ 544.2
93	-SO ₂ NH ₂	-H	-O-CH ₂ CH ₂ -OH	-OCH ₃	-H		
94	-OH	-H	-O-(1-methyl)-azacyclopent-2-yl	-H	-H		
95	-OH	-H	-O-CH ₂ CH ₂ -1-imidazolyl	-H	-H	389	387
96	-OH	-H	-O-CH ₂ CH ₂ CH ₂ -1-imidazolyl	-OCH ₃	-H	433.4	431.4
97	-SO ₂ NH ₂	-H	-OCH ₃	-H	-OCH ₃		
98	-OH	-H	-OCH ₃	-OCH ₃	-H	339	337
99	-SO ₂ NHCH ₂ -CH ₂ CH ₂ CH ₃	-H	-OCH ₃	-OCH ₃	-OCH ₃	488	486
100	-SO ₂ NH-CH ₃	-CH ₃	-O-CH ₂ CH ₂ -1-imidazolyl	-OCH ₃	-H	510	508
101	-SO ₂ NHCH ₂ -CH ₂ CH ₂ CH ₃	-H	-O-CH ₂ CH ₂ -1-imidazolyl	-H	-H	08	506
102	-OH	-H	-O-CH ₂ CH ₂ -4-morpholino	-H	-H	408	
103	-OH	-H	-NH-N=CH-		-H	319	317
104	-OH	-H	-CHN-NH-		-H	319	317
105	-OH	-H	-O-CH ₂ CH ₂ -1-imidazolyl	-H	-H		

106	-SO ₂ NH-CH ₃	-CH ₂ -CH ₃	-OCH ₃	-OCH ₃	-OCH ₃	474.3	472.3
107	-SO ₂ NH ₂	-H	-OCH ₃	-OCH ₃	-OCH ₃		

The compounds of formula X₅




wherein R⁰, R¹, R², R³ and R⁴ are as defined in Table 5, may be prepared by following the procedure of Example 1 but using the appropriate starting materials.

TABLE 5

Example	R ⁰	R ¹	R ²	R ³	R ⁴	MS Data	
						*ES+	*ES-
108	-H	-OCH ₃	-OH	-H	-H		
109	-H	nitro	-H	-OH	-H	414	412
110	-H	-N=CH-CH=CH-	-H	-H	-H		
111	-H	-CH=N-NH-	-H	-H	-H	393	391
112	-H	-NH-N=CH-	-H	-H	-H	393	
113	-H	-H	-OH	-CH ₂ CH ₂ CH ₂ -	-H	409	407
114	-CH ₃	-H	-CH ₃	-OH	-H	397	
115	-H	phenyl	-H	-SO ₂ NH ₂	-H	508	506
116	-CH ₃	-H	-H	-SO ₂ NH ₂	-H	446	444

The compounds of formula X₆




wherein R⁵, R⁷, R⁸ and R⁹ are as defined in Table 6, may be prepared by following the procedure of Example 1 but using the appropriate starting materials.

TABLE 6

Example	R ⁵	R ⁷	R ⁸	R ⁹	*ES+	*ES-
117	-CH ₃	-O-CH ₂ CH ₂ -1-imidazolyl	-H	-H	466	
118	-CH ₂ CH ₃	-OCH ₃	-OCH ₃	-OCH ₃	460	458
119	-Br	-NH-N=CH-		-H	461	
120	-CH ₃	-O-CH ₂ CH ₂ -1-imidazolyl	-OCH ₃	-H	496	
121	-CH ₃	-OCH ₃	-OCH ₃	-OCH ₃	446	
122	-CH ₃	-N=N-NH-		-H	397.2	395.2
123	-CH ₃	-O-CH ₂ CH ₂ -1-methyl-imidazol-1-yl	-H	-H	480	
124	-Br	-CH=N-NH-		-H	461.3	458.1 /460
125	-CH ₃	-NH-N=CH-		-H	396	
126	-Br	-OCH ₂ CH ₂ -(4-methyl-piperazin-1-yl)	-H	-H	562/ 564	560/ 562

The compounds of formula X₇




wherein R¹, R², R³, R⁷ and R⁸ are as defined in Table 7, may be prepared by following the procedure of Example 1 but using the appropriate starting materials.

TABLE 7

Ex	R ¹	R ²	R ³	R ⁷	R ⁸	*ES+	*ES-
127	-OCH ₃	-OH	-H	-OH	-OCH ₃		
128	-H	-CH ₃	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -1-imidazolyl	-H	466	464
129	-OCH ₃	-OH	-H	-O-CH ₂ CH ₂ -1-imidazolyl	-OCH ₃		
130	-OCH ₃	-OH	-H	-O-CH ₂ CH ₂ -OH	-OCH ₃	399	397
131	-OCH ₃	-OH	-H	-O-(1-methyl-azacyclohept-4-yl)	-H	436	
132	-CH ₃	-H	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -1-imidazolyl	-H	466	464
133	-OCH ₃	-OH	-H	-O-CH ₂ CH ₂ -(1-methyl)-azacyclopent-2-yl	-H	436	434
134	-OCH ₃	-OH	-H	-CF ₃	-H		
135	-N=CH-CH=CH-		-H	-O-CH ₂ CH ₂ -1-imidazolyl	-OCH ₃		
136	-OCH ₃	-OH	-H	-O-CH ₂ CH ₂ CH ₂ -1-imidazolyl	-OCH ₃	463	461
137	-OCH ₃	-OH	-H	-O-CH ₂ CH ₂ -1-piperidyl	-OCH ₃	466.4	464.4
138	-CH=N-NH-		-H	-NH-N=CH-			
139	-CH=N-NH-		-H	-CH=N-NH-			
140	-OCH ₃	-OH	-H	-O-CH ₂ CH ₂ -1-piperidyl	-H	436	434
141	-H	-OCH ₃	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -1-pyrrolidinyl	-H	485.3	483.3
142	-H	-OCH ₃	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -1-pyrrolidinyl	-CH ₃	499.2	497.3
143	-H	-OCH ₃	-SO ₂ NH ₂	-O-CH ₂ CH ₂ CH ₂ -morpholino	-OCH ₃	545.2	545.3

144	-H	-OCH(CH ₃) ₂	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -(4-methyl-piperazin-1-yl)	-OCH ₃	572.2	570.3
145	-H	-OCH ₃	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -1-piperidinyl	-H	499.2	497.3
146	-CH ₃	-OCH ₃	-SO ₂ NH ₂	-O-CH ₂ CH ₂ CH ₂ -1-pyrrolidinyl	-OCH ₃	543.2	
147	-CH ₃	-OCH ₃	-SO ₂ NH ₂	-O-CH ₂ CH ₂ CH ₂ -1-pyrrolidinyl	-H	513.2	511.2
148	-H	-OCH(CH ₃) ₂	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -1-piperidinyl	-H	527.2	525.3
149	-H	-CH ₃	-SO ₂ NH ₂	-N(CH ₃) ₂	-OCH ₃	429.3	427.3
150	-CH ₃	-CH ₃	-SO ₂ NH ₂	-O-CH ₂ CH ₂ CH ₂ -1-pyrrolidinyl	-OCH ₃	527.2	525.3
151	-OCH ₃	-H	-SO ₂ NH ₂	-O-CH ₂ CH ₂ CH ₂ -1-pyrrolidinyl	-OCH ₃	529.2	527.3
152	-H	-F	-SO ₂ NH ₂	-N(CH ₃) ₂	-OCH ₃	433.1	
153	-H	-CH ₃	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -(1-methyl-pyrrolidin-2-yl)	-H		
154	-H	-OCH ₃	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -OH	-H	432.2	430.2
155	-H	-CH ₃	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -(1-methyl-pyrrolidin-2-yl)	-OCH ₃	513.2	511.3
156	-OCH ₃	-H	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -1-piperidinyl	-H	499.2	497.3
157	-OCH ₃	-H	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -1-pyrrolidinyl	-OCH ₃	515.2	513.2
158	-H	-CH ₃	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -OH	-OCH ₃	446.2	444.2
159	-OC ₂ H ₅	-H	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -1-pyrrolidinyl	-CH ₃	513.3	511.3
160	-OCH ₃	-OCH ₃	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -(4-methyl-piperazin-1-yl)	-OCH ₃	574.2	572.2
161	-H	-Cl	-SO ₂ NH ₂	-(4-methyl-piperazin-1-yl)	-H	474.5	472.5
162	-H	-CH ₃	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -(4-cyclopentyl-piperazin-1-yl)	-H	552.3	550.3
163	-CH=CH-CH=CH-		-SO ₂ NH ₂	-(4-methyl-piperazin-1-yl)	-H	490.5	488.4
164	-H	-H	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -piperazin-1-yl	-H	470.2	468.3
165	-H	-OCH ₃	-SO ₂ NH ₂	-H	-OCH ₃	402.2	400.2
166	-H	-OCH ₃	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -(4-benzyl-piperazin-1-yl)	-H	590.3	588.3
167	-CH ₃	-H	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -1-pyrrolidinyl	-H	469.2	467.3
168	-Br	-H	-SO ₂ NH ₂	-O-CH ₂ CH ₂ -1-piperidinyl	-H	549.1	547.2

The compounds of formula X₈




wherein R^1 , R^2 , R^3 and R^8 are as defined in Table 8, may be prepared by following the procedure of Example 1 but using the appropriate starting materials.

TABLE 8

Ex	R^1	R^2	R^3	R^8	*ES+	*ES-
169	4-morpholino	-H	-H	-H		
170		-CH=N-NH-	-H	-H	363	361
171	-OCH ₃	-OH	-H	-H		
172	-CH ₃	-H	-SO ₂ NH ₂	-OCH ₃	446	

The compounds of formula X₉




wherein R^7 , R^8 and R^9 are as defined in Table 9, may be prepared by following the procedure of Example 1 but using the appropriate starting materials.

TABLE 9

Example	R^1	R^8	R^9	*ES+	*ES-
173	-O-CH ₂ CH ₂ -1-piperidyl	-OCH ₃	-H	470.3	468.3
174	-O-(1-methyl-azacyclohept-4-yl)	-H	-H	440	
175	-O-(1-methyl-azacyclopent-2-yl)	-H	-H	440	438
176	-O-CH ₂ CH ₂ -CH ₂ -1-imidazolyl	-OCH ₃	-H	467	465
177	-OCH ₃	-OCH ₃	-OCH ₃		
178	-O-CH ₂ CH ₂ -1-(1,2,4-triazolyl)	-H	-H	424	422
179	-O-CH ₂ CH ₂ -1-piperidyl	-H	-H		
180	-O-CH ₂ CH ₂ -OH	-OCH ₃	-H		
181	-O-CH ₂ CH ₂ -4-morpholino	-H	-H	442	440
182	-O-CH ₂ CH ₂ -CH ₂ -1-imidazolyl	-H	-H		

The compounds of formula X₁₀




wherein R^1 , R^7 and R^9 are as defined in Table 10, may be prepared by following the procedure of Example 1 but using the appropriate starting materials.

TABLE 10


EX	R^1	R^7	R^9	*ES+	*ES-
183	$-\text{CH}_2\text{CH}_2-\text{OH}$	$-\text{OCH}_3$	$-\text{OCH}_3$	411	409
184	$-\text{SO}_2\text{NH}_2$	$-\text{O}-\text{CH}_2\text{CH}_2-1\text{-imidazolyl}$	$-\text{H}$	496.3	494.3

The compounds of formula X_{11}



wherein R^8 is $-\text{OCH}_3$ (Example 185) or $-\text{OH}$ (Example 186), may be prepared by following the procedure of Example 1 but using the appropriate starting materials.

The compounds of formula X_{12}




wherein R^0 , R^1 , R^7 , R^8 and R^9 are as defined in Table 12, may be prepared by following the procedure of Example 1 but using the appropriate starting materials.

TABLE 12

Example	R^0	R^1	R^7	R^8	R^9
187	$-\text{H}$	$-\text{H}$	$-\text{H}$	$-\text{SO}_2\text{NH}_2$	$-\text{H}$
188	$-\text{H}$	$-\text{H}$	$-\text{H}$	$-\text{H}$	$-\text{CH}_3$
189	$-\text{H}$	$-\text{H}$	$-\text{H}$	$-\text{CH}_3$	$-\text{H}$
190	$-\text{H}$	$-\text{F}$	$-\text{OCH}_3$	$-\text{OCH}_3$	$-\text{OCH}_3$
191	$-\text{H}$	$-\text{H}$	$-\text{H}$	$-\text{CH}_3$	$-\text{CH}_3$
192	$-\text{H}$	$-\text{H}$	$-\text{CH}_3$	$-\text{H}$	$-\text{CH}_3$
193	$-\text{H}$	$-\text{H}$	$-\text{OCH}_3$	$-\text{CH}_3$	$-\text{H}$
194	$-\text{H}$	$-\text{H}$	$-\text{H}$	$-\text{H}$	$-\text{N}(\text{CH}_3)_2$

195	-H	-H	-OCH(CH ₃) ₂	-H	-H
196	-H	-H	-H	-OCH(CH ₃) ₂	-H
197	-H	-H	-CH(CH ₃) ₂	-H	-H
198	-H	-H	-H	-CH=N-NH-	
199	-H	-H	-OCH ₃	-CH ₃	-OCH ₃
200	-OCH ₃	-H	-OCH ₃	-OCH ₃	-OCH ₃
201	-H	-H	-H	-H	-H
202	-CH ₃	-Cl	-OCH ₃	-OCH ₃	-OCH ₃
203	-H	-H	-H	-H	-CF ₃
204	-Cl	-CH ₃	-OCH ₃	-OCH ₃	-OCH ₃
205	-H	-H	-H	-NH-CH=N-	
206	-H	-H	-H	-N(-CH ₂ CH ₂ CH ₂ -4-morpholino)-CH=CH-	
207	-H	-H	-CH ₂ CH ₂ -CH ₂ -		-H

The compounds of formula X₁₃




wherein R¹, R², R³ and R⁵ are as defined in Table 13, may be prepared by following the procedure of Example 1 but using the appropriate starting materials.

TABLE 13

Example	R ¹	R ²	R ³	R ⁵	*ES+	*ES-
208	-H	-H	-SO ₂ NHCH ₃	-CF ₃	514.0	
209	-H	-H	-SO ₂ NHC ₃ H ₇	-Br		
210	-H	-H	-SO ₂ NH-CH ₂ CH-cyclopropyl	-Br		
211	-H	-H	-SO ₂ NHCH ₃	-CH ₃		
212	-H	-H	-SO ₂ N(CH ₃) ₂	-Br		
213	-H	-H	-SO ₂ NHCH ₃	-Cl		
214	-H	-H	-SO ₂ NHCH ₃	-I		
215	-H	-H	-SO ₂ NHCH ₃	-Br		
216	-CH ₃	-OCH ₃	-SO ₂ NH ₂	-H	476	474
217	-H	piperidino	-SO ₂ NH ₂	-H	515.5	513.4
218	-H	morpholino	-SO ₂ NH ₂	-H	517.4	515.4

219	-H	-C ₂ H ₅	-SO ₂ NH ₂	-H		
220	-H	-CH ₃	-SO ₂ NH ₂	-Cl		
221	-H	-CH ₃	-SO ₂ NHCH ₃	-H	460.4	
222	-H	phenyl	-SO ₂ NH ₂	-H	508.2	506.3

The compounds of formula X₁₄



wherein R², R³, R⁵, R⁷, R⁸ and R⁹ are as defined in Table 14, may be prepared by following the procedure of Example 1 but using the appropriate starting materials.

TABLE 14

Ex	R ²	R ³	R ⁵	R ⁷	R ⁸	R ⁹	*ES+	*ES-
223	-OCH ₃	-SO ₂ NH ₂	-H	-H	-CH=N-N(CH ₃)-			424
224	-OCH ₃	-SO ₂ NH ₂	-H	-O-CH ₂ CH ₂ -OCH ₃	-OCH ₃	-H	476.2	474.3
225	-OCH(CH ₃) ₂	-SO ₂ NH ₂	-H	-O-CH ₂ CH ₂ - piperidino	-OCH ₃	-H	551.2	555.3
226	-OCH ₃	-SO ₂ NH ₂	-H	-O-CH ₂ CH ₂ -(4-methyl-piperazin-1-yl)	-H	-H	514.3	512.3
227	-OCH ₃	-SO ₂ NH ₂	-H	-morpholino	-OCH ₃	-H	487.1	485.2
228	-CH ₃	-SO ₂ NH ₂	-H	-O-CH ₂ CH ₂ CH ₂ - piperidino	-OCH ₃	-H	527.3	
229	-CH ₃	-SO ₂ NH ₂	-H	-O-CH ₂ CH ₂ CH ₂ -1-pyrrolidinyl	-OCH ₃	-H	513.2	511.3
230	-O-CH ₂ CH ₂ -OCH ₃	-SO ₂ NH ₂	-H	-H	-CH=N-N(CH ₃)-		539	537
231	-(4-methyl-piperazin-1-yl)	-SO ₂ NH ₂	-H	-OCH ₃	-OCH ₃	- OCH ₃	530.4	528.4
232	-OCH ₃	-SO ₂ NH ₂	-H	-O-CH ₂ CH ₂ -OH	-OCH ₃	-H	462.2	460.3
233	-OCH ₃	-SO ₂ NH ₂	-Br	-O-CH ₂ CH ₂ -OCH ₃	-OCH ₃	-H		
234	-CH ₃	-SO ₂ NH ₂	-H	-O-CH ₂ CH ₂ -(4-methyl-piperazin-1-yl)	-OCH ₃	-H	528.2	526.3
235	-CH ₃	-SO ₂ NH ₂	-H	-O-CH ₂ CH ₂ -N(CH ₃) ₂	-H	-H	443.2	441.3
236	-H	-SO ₂ NH ₂	-H	-O-CH ₂ CH ₂ -1-pyrrolidinyl	-OCH ₃	-H	485.2	483.3
237	-CH ₃	-SO ₂ NH ₂	-H	-H	-N(CH ₃)-N=CH-		410	
238	-CH ₃	-SO ₂ NH ₂	-H	-CH ₃	-OCH ₃	OCH ₃		

239	-CH ₃	-SO ₂ NH ₂	-Br	-O-CH ₂ CH ₂ -OCH ₃	-OCH ₃	-H	538/ 540	
240	-OCH ₃	-SO ₂ NH ₂	-H	-OCH ₃	-H	-H	402.2	400.2
241	-H	-SO ₂ NH ₂	-H	-H	-CO- NH- CH ₂ CH ₂ - OCH ₃	-H		

ES+ means electrospray MS positive mode ; ES- means electrospray MS negative mode; and EL means electron impact MS.

The compounds of formula I and their pharmaceutically acceptable salts, exhibit valuable pharmacological properties when tested in in vitro assays, and are therefore useful as pharmaceuticals.

In particular the compounds of the invention exhibit ZAP-70 (zeta chain-associated protein of 70 kD), Focal Adhesion Kinase (FAK) and/or Syk protein tyrosine kinases inhibiting activity. More particularly the compounds of the invention are active at the human ZAP-70, FAK and/or Syk protein tyrosine kinases. ZAP-70, FAK and/or Syk protein tyrosine kinase interaction of the compounds of the invention may be demonstrated by their ability to prevent phosphorylation of e.g. LAT-11 (SEQ ID NO: 1) by human ZAP-70 protein tyrosine kinase, to prevent phosphorylation of e.g. Biot-Y397 (SEQ ID NO:2) by human FAK protein tyrosine kinase, and/or to prevent phosphorylation of e.g. polymeric glutamic acid-tyrosine (Glu, Tyr) by human Syk protein tyrosine kinase in, e.g. aqueous solution, e.g. as demonstrated in accordance with the following test methods.

1. Cell-free kinase assays: ZAP-70 and Syk Kinase assays

ZAP-70, Lck and Syk are commercially available from Upstate Biotechnology, Lake Placid, NY.

Preparation of LAT-11 (SEQ ID NO:1): The peptide LAT-11 used as a substrate in the ZAP-70 kinase assay may be prepared as disclosed in Example 1A of WO 02/12275, the contents of which, particularly with reference to Example 1A, is incorporated herein by reference.

ZAP-70 Kinase assay: The activities of the agents of invention are determined in a homogenous ZAP-70 kinase assay based on time-resolved fluorescence resonance energy transfer. Briefly, 80 nM ZAP-70 are incubated with 80 nM Lck and 4 μ M ATP in ZAP-70 kinase buffer (20 mM Tris, pH 7.5, 10 μ M Na₃VO₄, 1 mM DTT, 1 mM MnCl₂, 0.01 % bovine serum albumin, 0.05 %

Tween 20) for 1 hour at room temperature in a siliconized polypropylene tube. Then, the selective Lck inhibitor PP2 (4-amino-5-(4-chloro-phenyl)-7-(t-butyl)pyrazolo[3,4-d]pyrimidine; Alexis Biochemicals) is added (final concentration 1.2 μM) and incubated for further 10 min. Ten μl of this solution is mixed with the 10 μl biotinylated peptide LAT-11 (1 μM) as substrate and 20 μl of serial dilutions of inhibitors and incubated for 4 hours at room temperature. The kinase reaction is terminated with 10 μl of a 10 mM EDTA solution in detection buffer (20 mM Tris, pH 7.5, 0.01 % bovine serum albumin, 0.05 % Tween 20). The detection phase is performed by addition of 50 μl europium (Eu)-labelled anti-phosphotyrosine antibody (e.g. Eu-PT66; final concentration 0.125 nM; Advant/Wallac) and 50 μl streptavidin-allophycocyanine (SA-APC; final concentration 40 nM) in detection buffer. After 1 hour incubation at room temperature fluorescence is measured, e.g., on the Victor2 Multilabel Counter (Wallac) at 665 nm. Background values (low control) are obtained in the absence of test samples and ATP and are subtracted from all values. Signals obtained in the absence of test samples are taken as 100% (high control). The inhibition obtained in the presence of test compounds was calculated as percent inhibition of the high control. The concentration of test compounds resulting in 50% inhibition (IC_{50}) was determined from the dose-response curves. In this assay, the compounds of the invention have IC_{50} values in the range of 10 nM to 2 μM , preferably from 10 nM to 100 nM. Compound of Example 4 shows an IC_{50} value of 12 nM.

Syk Kinase assay: The activities of the agents of invention are determined in a heterogeneous Syk kinase assay based on the dissociation-enhanced lanthanide fluoroimmunoassay (DELFI) technology. This method utilizes europium chelate-labelled anti-phosphotyrosine antibodies to detect phosphate transfer by Syk to a polymeric glutamic acid-tyrosine (Glu, Tyr) substrate coated onto microtiter plates as described (Braunwalder AF, Yarwood DR, Sills MA, Lipson KE. Measurement of the protein tyrosine kinase activity of c-src using time-resolved fluorometry of europium chelates. *Anal.Biochem.* 1996;238(2):159-64). The amount of phosphorylation is then quantified with time-resolved, dissociation-enhanced fluorescence. Briefly, hundred μl of poly (Glu, Tyr) (4:1; 2 $\mu\text{g/ml}$ in phosphate-buffered saline, PBS) are coated to ELISA plates overnight at room temperature. The poly (Glu, Tyr) solution is removed and 250 μl of 1% bovine serum albumin in PBS are added for one hour at room temperature. Plates are then washed three times with 350 μl of washing buffer (25 mM Tris-HCl, pH 7.4 containing 0.03% Tween-20). The kinase reaction is performed for one hour at room temperature by mixing serial dilutions of inhibitors in 30 μl with 30 μl of Syk kinase (20 ng/ml) and ATP (1 μM) in kinase buffer (20 mM Tris, pH 7.5, 10 μM Na_3VO_4 , 1 mM DTT, 10 mM MnCl_2 , 2 mM MgCl_2 , 0.01 % bovine serum albumin, 0.05 % Tween 20). After

washing the plates four times as described above 60 μ l DELFIA europium N1-labelled anti-phosphotyrosine antibody PY20 (Advant/Wallac) are added (100 ng/ml in 50 mM Tris-HCl, pH7.4, 150 mM NaCl, 20 μ M Titriplex V, 0.2% bovine serum albumine, 0.05% Tween-20) and incubated for one hour at room temperature. Plates are washed eight times and 60 μ l enhancement solution (Wallac) are added. Fluorescence is determined at 615 nm (Victor2; Wallac). High control values (100% signal) are obtained in absence of test samples and low control values (background) in absence of test samples and ATP. Low controls were subtracted from all values. The inhibition obtained in the presence of test compounds was calculated as percent inhibition of the high control. The concentration of test compounds resulting in 50% inhibition (IC_{50}) was determined from the dose-response curves. In this assay, the compounds of the invention have IC_{50} values in the range of 100 nM to 10 μ M, preferably from 100 to 1 μ M. Compound of Example 128 has an IC_{50} value of 150 nM.

2. Allogeneic Mixed Lymphocyte Reaction (MLR)

Compounds of the invention exhibit T cell inhibiting activity. More particular the compounds of the invention prevent T cell activation and/or proliferation in e.g. aqueous solution, e.g. as demonstrated in accordance with the following test method. The two-way MLR is performed according to standard procedures (J. Immunol. Methods, 1973, 2, 279 and Meo T. et al., Immunological Methods, New York, Academic Press, 1979, 227-39). Briefly, spleen cells from CBA and BALB/c mice (1.6×10^5 cells from each strain per well in flat bottom tissue culture microtiter plates, 3.2×10^5 in total) are incubated in RPMI medium containing 10% FCS, 100 U/ml penicillin, 100 μ g/ml streptomycin (Gibco BRL, Basel, Switzerland), 50 μ M 2-mercaptoethanol (Fluka, Buchs, Switzerland) and serially diluted compounds. Seven three-fold dilution steps in duplicates per test compound are performed. After four days of incubation 1 μ Ci 3 H-thymidine is added. Cells are harvested after an additional five-hour incubation period, and incorporated 3 H-thymidine is determined according to standard procedures. Background values (low control) of the MLR are the proliferation of BALB/c cells alone. Low controls are subtracted from all values. High controls without any sample are taken as 100% proliferation. Percent inhibition by the samples is calculated, and the concentrations required for 50% inhibition (IC_{50} values) are determined. In this assay, the compounds of the invention have IC_{50} values in the range of 10 nM to 10 μ M, preferably from 10 nM to 100 nM. Compound of Example 120 shows an IC_{50} value of 13 nM.

3. FAK assay

All steps are performed in a 96-well black microtiter plate. Purified recombinant hexahistidine-tagged human FAK kinase domain is diluted with dilution buffer (50 mM HEPES, pH 7.5, 0.01% BSA, 0.05% Tween-20 in water) to a concentration of 94 ng/mL (2.5 nM). The reaction mixture is prepared by mixing 10 μ L 5x kinase buffer (250 mM HEPES, pH 7.5, 50 μ M Na₃VO₄, 5 mM DTT, 10 mM MgCl₂, 50 mM MnCl₂, 0.05% BSA, 0.25% Tween-20 in water), 20 μ L water, 5 μ L of 4 μ M biotinylated peptide substrate (Biot-Y397) in aqueous solution, 5 μ L of test compound in DMSO, and 5 μ L of recombinant enzyme solution and incubated for 30 min at room temperature. The enzyme reaction is started by addition of 5 μ L of 5 μ M ATP in water and the mixture is incubated for 3 hours at 37°C. The reaction is terminated by addition of 200 μ L of detection mixture (1 nM Eu-PT66, 2.5 μ g/mL SA-(SL)APC, 6.25 mM EDTA in dilution buffer), and the FRET signal from europium to allophycocyanin is measured by ARVOSx+L (Perkin Elmer) after 30 min of incubation at room temperature. The ratio of fluorescence intensity of 665 nm to 615 nm is used as a FRET signal for data analysis in order to cancel the colour quenching effect by a test compound. The results are determined as percent inhibition of enzyme activity. DMSO and 0.5 M EDTA are used as a control of 0% and 100% inhibition, respectively. IC₅₀ values are determined by non-linear curve fit analysis using the OriginPro 6.1 program (OriginLab). In this assay the compounds of formula I inhibit FAK activity at a IC₅₀ < 1 μ M. Examples 188, 208 and 213 show IC₅₀ values of 15 nM, 1 nM and 7 nM respectively.

The Biot-Y397 peptide (Biotin-SETDDYAEIID ammonium salt, SEQ ID NO:2) is designed to have the same amino acid sequence as the region from S392 to D402 of human (GenBank Accession Number L13616) and is prepared by standard methods.

Purified recombinant hexahistidine-tagged human FAK kinase domain is obtained in the following way: Full-length human FAK cDNA is isolated by PCR amplification from human placenta Marathon-Ready™ cDNA (Clontech, No. 7411-1) with the 5' PCR primer (ATGGCAGCTGCTTACCTTGAC, SEQ ID NO:3) and the 3' PCR primer (TCAGTGTGGTCTCGTCTGCCC, SEQ ID NO:4) and subcloned into a pGEM-T vector (Promega, No. A3600). After digestion with AccIII, the purified DNA fragment is treated with Klenow fragment. The cDNA fragment is digested with BamHI and cloned into pFastBachTb plasmid (Invitrogen Japan K.K., Tokyo) previously cut with BamHI and Stu I. The resultant plasmid, hFAK KD (M384-G706)/pFastBachTb, is sequenced to confirm its structure. The resulting DNA encodes a 364 amino acid protein containing a hexahistidine tag, a spacer region

and a rTEV protease cleavage site at the N-terminal and the kinase domain of FAK (Met384-Gly706) from position 29 to 351.

Donor plasmid is transposed into the baculovirus genome, using MaxEfficacy DH10Bac *E.coli* cells. Bacmid DNA is prepared by a simple alkaline lysis protocol described in the Bac-to-Bac® Baculovirus Expression system (Invitrogen). Sf9 insect cells are transfected based on the protocol provided by the vendor (CellFECTIN®, Invitrogen). The expression of FAK in each lysate is analysed by SDS-PAGE and Western blotting with anti-human FAK monoclonal antibody (clone #77 from Transduction Laboratories).

The virus clone that shows the highest expression is further amplified by infection to Sf9 cells. Expression in ExpressSF+® cells (Protein Sciences Corp., Meriden, Connecticut, USA) gives high level of protein with little degradation. Cell lysates are loaded onto a column of HiTrap™ Chelating Sepharose HP (Amersham Biosciences) charged with nickel sulfate and equilibrated with 50 mM HEPES pH 7.5, 0.5 M NaCl and 10 mM imidazole. Captured protein is eluted with increasing amounts of imidazole in HEPES buffer / NaCl, and further purified by dialysis in 50 mM HEPES pH 7.5, 10% glycerol and 1 mM DTT.

4. Phosphorylation levels of FAK

Phosphorylation levels of FAK at Tyr397 are quantified by the sandwich ELISA. Mouse mammary carcinoma 4T1 cells (1×10^5) are plated in wells of 96-well culture plates and incubated with or without various concentrations of a compound of formula I for 1 h in Dulbecco's modified eagle medium containing 10% FBS. The medium is removed and cells are lysed in 200 μ L 50 mM Tris-HCl, pH 7.4, containing 1% NP-40, 0.25% sodium deoxycholate, 150 mM NaCl, 1 mM EDTA, 1 mM PMSF, 1 mM Na_3VO_4 , 1 mM NaF, 1 μ g/mL aprotinin, 1 μ g/mL leupeptin and 1 μ g/mL pepstatin. After centrifugation, the supernatants are subjected to a sandwich ELISA to quantify the phosphorylated FAK and total FAK. Cell lysates are applied to 96-well flat-bottom ELISA plates which have been pre-coated with 100 μ L/well of 4 μ g/mL mouse monoclonal anti-FAK antibody (clone 77, Becton Dickinson Transduction Laboratories) in 50 mM Tris-HCl, pH 9.5, containing 150 mM NaCl for 18 h at 4°C and blocked with 300 μ L of BlockAce (Dainippon Pharmaceuticals Co.) diluted at 1:4 with H_2O at room temperature for 2 h. After washing with TBSN (20 mM Tris-HCl, pH 8.3, containing 300 mM NaCl, 0.1% SDS and 0.05% NP-40), total FAK is detected with 100 μ L of 1 μ g/ml anti-FAK polyclonal antibody (#65-6140, Upstate Biology Inc.), and phosphorylated FAK is detected with 100 μ L of 0.25 μ g/ μ L anti-

phosphorylated FAK (Y397) antibody (Affinity BioReagents, #OPA1-03071) in BlockAce diluted at 1:10 with H₂O. After 1 h incubation at room temperature, plates are washed with TBSN and 100 μ L of biotinylated anti-rabbit IgG (#65-6140, Zymed Laboratories Inc.) diluted at 1:2000 with BlockAce diluted at 1:10 with H₂O is incubated at room temperature for 1 h. After washing with TBSN, ABTS solution substrate kit (#00-2011, Zymed Laboratories Inc.) is used for color development. Absorbance at 405 nm is measured after 20 min incubation at room temperature. The concentration of compound causing 50% reduction of phosphorylation level of FAK (IC₅₀) is determined. In this assay, compounds of formula I reduce phosphorylation at an IC₅₀ of < 1 μ M. Examples 190, 198 and 210 show IC₅₀ values of 0.44 μ M, 0.043 μ M and 0.01 μ M respectively.

5. Anchorage-independent tumor cell growth assay

Mouse mammary carcinoma 4T1 cells (5×10^3) are plated in 96-well Ultra low Attachment plates (#3474, Corning Inc.) in 100 μ L of Dulbecco's modified eagle medium containing 10% FBS. Cells are cultured for 2 h and inhibitors are added at various concentrations in a final concentration of 0.1% DMSO. After 48 h, cell growth is assayed with the cell counting kit-8 (Wako Pure Chemical), which uses a water soluble tetrazolium salt WST8. Twenty μ L of the reagent is added into each well and cells are further cultured for 2 h. The optical density is measured at 450 nm. The concentration of compound causing 50 % inhibition of growth may thus be determined. Examples 204, 213 and 206 show IC₅₀ values of 0.4 μ M, 0.016 μ M and 0.09 μ M respectively.

The compounds of the invention are therefore useful in the prevention or treatment of disorders or diseases where ZAP-70 inhibition, and/or Syk inhibition play a role, e.g. diseases or disorders mediated by T lymphocytes, B lymphocytes, mast cells and/or eosinophils e.g. acute or chronic rejection of organ or tissue allo- or xenografts, atherosclerosis, vascular occlusion due to vascular injury such as angioplasty, restenosis, hypertension, heart failure, chronic obstructive pulmonary disease, CNS disease such as Alzheimer disease or amyotrophic lateral sclerosis, cancer, infectious disease such as AIDS, septic shock or adult respiratory distress syndrome, ischemia/reperfusion injury e.g. myocardial infarction, stroke, gut ischemia, renal ailure or hemorrhage shock, or traumatic shock. The agent of the invention are also useful in the treatment and/or prevention of acute or chronic inflammatory diseases or disorders or autoimmune diseases e.g. rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, Hashimoto's thyroidis, multiple sclerosis, myasthenia gravis, diabetes (type I and II) and the disorders associated with therewith, respiratory diseases such as asthma or inflammatory liver

injury, inflammatory glomerular injury, cutaneous manifestations of immunologically-mediated disorders or illnesses, inflammatory and hyperproliferative skin diseases (such as psoriasis, atopic dermatitis, allergic contact dermatitis, irritant contact dermatitis and further eczematous dermatitises, seborrhoeic dermatitis), inflammatory eye diseases, e.g. Sjogren's syndrome, keratoconjunctivitis or uveitis, inflammatory bowel disease, Crohn's disease or ulcerative colitis.

Compounds of the invention are also useful in the prevention or treatment of conditions caused by a malfunction of signal cascades connected with FAK, e.g. tumors, for example brain and other central nervous system tumors (eg. tumors of the meninges, brain, spinal cord, cranial nerves and other parts of central nervous system, e.g. glioblastomas or medulla blastomas); head and/or neck cancer; breast tumors; circulatory system tumors (e.g. heart, mediastinum and pleura, and other intrathoracic organs, vascular tumors and tumor-associated vascular tissue); excretory system tumors (e.g. kidney, renal pelvis, ureter, bladder, other and unspecified urinary organs); gastrointestinal tract tumors (e.g. oesophagus, stomach, small intestine, colon, colorectal, rectosigmoid junction, rectum, anus and anal canal), tumors involving the liver and intrahepatic bile ducts, gall bladder, other and unspecified parts of biliary tract, pancreas, other and digestive organs); head and neck; oral cavity (lip, tongue, gum, floor of mouth, palate, and other parts of mouth, parotid gland, and other parts of the salivary glands, tonsil, oropharynx, nasopharynx, pyriform sinus, hypopharynx, and other sites in the lip, oral cavity and pharynx); reproductive system tumors (e.g. vulva, vagina, Cervix uteri, Corpus uteri, uterus, ovary, and other sites associated with female genital organs, placenta, penis, prostate, testis, and other sites associated with male genital organs); respiratory tract tumors (e.g. nasal cavity and middle ear, accessory sinuses, larynx, trachea, bronchus and lung, e.g. small cell lung cancer or non-small cell lung cancer); skeletal system tumors (e.g. bone and articular cartilage of limbs, bone articular cartilage and other sites); skin tumors (e.g. malignant melanoma of the skin, non-melanoma skin cancer, basal cell carcinoma of skin, squamous cell carcinoma of skin, mesothelioma, Kaposi's sarcoma); and tumors involving other tissues including peripheral nerves and autonomic nervous system, connective and soft tissue, retroperitoneum and peritoneum, eye and adnexa, thyroid, adrenal gland and other endocrine glands and related structures, secondary and unspecified malignant neoplasm of lymph nodes, secondary malignant neoplasm of respiratory and digestive systems and secondary malignant neoplasm of other sites, tumors of blood and lymphatic system (e.g. Hodgkin's disease, Non-Hodgkin's lymphoma, Burkitt's lymphoma, AIDS-related lymphomas, malignant immunoproliferative diseases, multiple myeloma and malignant plasma cell neoplasms, lymphoid leukemia, acute or chronic myeloid leukemia, acute or chronic lymphocytic leukemia,

monocytic leukemia, other leukemias of specified cell type, leukemia of unspecified cell type, other and unspecified malignant neoplasms of lymphoid, haematopoietic and related tissues, for example diffuse large cell lymphoma, T-cell lymphoma or cutaneous T-cell lymphoma). Myeloid cancer includes e.g. acute or chronic myeloid leukaemia.

Where hereinbefore and subsequently a tumor, a tumor disease, a carcinoma or a cancer is mentioned, also metastasis in the original organ or tissue and/or in any other location are implied alternatively or in addition, whatever the location of the tumor and/or metastasis is.

The compositions of the invention may be administered by any conventional route, in particular parenterally, for example in the form of injectable solutions or suspensions, enterally, e.g. orally, for example in the form of tablets or capsules, topically, e.g. in the form of lotions, gels, ointments or creams, or in a nasal or a suppository form. Pharmaceutical compositions comprising an agent of the invention in association with at least one pharmaceutical acceptable carrier or diluent may be manufactured in conventional manner by mixing with a pharmaceutically acceptable carrier or diluent. Unit dosage forms for oral administration contain, for example, from about 0.1 mg to about 500 mg of active substance. Topical administration is e.g. to the skin. A further form of topical administration is to the eye.

The compounds of formula I may be administered in free form or in pharmaceutically acceptable salt form, e.g. as indicated above. Such salts may be prepared in conventional manner and exhibit the same order of activity as the free compounds.

In accordance with the foregoing, the present invention also provides:

- (1) A compound of formula I or a pharmaceutically acceptable salt thereof, for use as a pharmaceutical;
- (2) A compound of formula I or a pharmaceutically acceptable salt thereof, for use as a ZAP-70, FAK and/or Syk tyrosine kinase inhibitor, for example for use in any of the particular indications hereinbefore set forth;

(3) A pharmaceutical composition, e.g. for use in any of the indications herein before set forth, comprising a compound of formula I or a pharmaceutically acceptable salt thereof, together with one or more pharmaceutically acceptable diluents or carriers therefor.

(4) A method for the treatment of any of particular indication hereinbefore set forth in a subject in need thereof which comprises administering an effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof;

(5) The use of a compound of formula I or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment or prevention of a disease or condition in which ZAP-70, FAK and/or Syk tyrosine kinase activation plays a role or is implicated; e.g. as discussed above.

Compounds of the invention may be administered as the sole active ingredient or together with other drugs useful against neoplastic diseases, inflammatory disorders or in immunomodulating regimens. For example, the compounds of the invention may be used in combination with an active agent effective in various diseases as described above, e.g. with cyclosporins, rapamycins or ascomycins, or their immunosuppressive analogs or derivatives, e.g. cyclosporin A, cyclosporin G, Isa tx247, FK-506, sirolimus or everolimus; CCI-779, ABT578, AP23573, corticosteroids e.g. prednisone; cyclophosphamide; azathioprene; methotrexate; gold salts, sulfasalazine, antimalarials; leflunomide; mizoribine; mycophenolic acid; mycophenolate mofetil; 15-deoxyspergualine; an EDG receptor agonist having accelerating lymphocyte homing activity, e.g. FTY720 or an analogue thereof, immuno-suppressive monoclonal antibodies, e.g. monoclonal antibodies to leukocyte receptors, e.g. MHC, CD2, CD3, CD4, CD7, CD25, CD28, CD40, CD45, CD58, CD80, CD86, CD152, CD137, CD154, ICOS, LFA-1, VLA-4 or their ligands; or other immunomodulatory compounds, e.g. CTLA4Ig.

A compound of formula I may also be used to advantage in combination with other antiproliferative agents. Such antiproliferative agents include, but are not limited to aromatase inhibitors, antiestrogens, topoisomerase I inhibitors, topoisomerase II inhibitors, microtubule active agents, alkylating agents, histone deacetylase inhibitors, farnesyl transferase inhibitors, COX-2 inhibitors, MMP inhibitors, mTOR inhibitors, antineoplastic antimetabolites, platinum compounds, compounds decreasing the protein kinase activity and further anti-angiogenic

compounds, gonadorelin agonists, anti-androgens, bengamides, bisphosphonates, antiproliferative antibodies and temozolomide (TEMODAL®).

The term "aromatase inhibitors" as used herein relates to compounds which inhibit the estrogen production, i.e. the conversion of the substrates androstenedione and testosterone to estrone and estradiol, respectively. The term includes, but is not limited to steroids, especially exemestane and formestane and, in particular, non-steroids, especially aminoglutethimide, vorozole, fadrozole, anastrozole and, very especially, letrozole. A combination of the invention comprising an antineoplastic agent which is an aromatase inhibitor may particularly be useful for the treatment of hormone receptor positive breast tumors.

The term "antiestrogens" as used herein relates to compounds which antagonize the effect of estrogens at the estrogen receptor level. The term includes, but is not limited to tamoxifen, fulvestrant, raloxifene and raloxifene hydrochloride.

The term "topoisomerase I inhibitors" as used herein includes, but is not limited to topotecan, irinotecan, 9-nitrocamptothecin and the macromolecular camptothecin conjugate PNU-166148 (compound A1 in WO99/17804).

The term "topoisomerase II inhibitors" as used herein includes, but is not limited to the anthracyclines doxorubicin (including liposomal formulation, e.g. CAELYX™), epirubicin, idarubicin and nemorubicin, the anthraquinones mitoxantrone and losoxantrone, and the podophyllotoxines etoposide and teniposide.

The term "microtubule active agents" relates to microtubule stabilizing and microtubule destabilizing agents including, but not limited to the taxanes paclitaxel and docetaxel, the vinca alkaloids, e.g., vinblastine, especially vinblastine sulfate, vincristine especially vincristine sulfate, and vinorelbine, discodermolide and epothilones, such as epothilone B and D.

The term "alkylating agents" as used herein includes, but is not limited to cyclophosphamide, ifosfamide and melphalan.

The term "histone deacetylase inhibitors" relates to compounds which inhibit the histone deacetylase and which possess antiproliferative activity.

The term "farnesyl transferase inhibitors" relates to compounds which inhibit the farnesyl transferase and which possess antiproliferative activity.

The term "COX-2 inhibitors" relates to compounds which inhibit the cyclooxygenase type 2 enzyme (COX-2) and which possess antiproliferative activity such as celecoxib (Celebrex®), rofecoxib (Vioxx®) and lumiracoxib (COX189).

The term "MMP inhibitors" relates to compounds which inhibit the matrix metalloproteinase (MMP) and which possess antiproliferative activity.

The term "antineoplastic antimetabolites" includes, but is not limited to 5-fluorouracil, tegafur, capecitabine, cladribine, cytarabine, fludarabine phosphate, fluorouridine, gemcitabine, 6-mercaptopurine, hydroxyurea, methotrexate, edatrexate and salts of such compounds, and furthermore ZD 1694 (RALTITREXED™), LY231514 (ALIMTA™), LY264618 (LOMOTREXOL™) and OGT719.

The term "platin compounds" as used herein includes, but is not limited to carboplatin, cis-platin and oxaliplatin.

The term "compounds decreasing the protein kinase activity and further anti-angiogenic compounds" as used herein includes, but is not limited to compounds which decrease the activity of e.g. the Vascular Endothelial Growth Factor (VEGF), the Epidermal Growth Factor (EGF), c-Src, protein kinase C, Platelet-derived Growth Factor (PDGF), Bcr-Abl tyrosine kinase, c-kit, Flt-3 and Insulin-like Growth Factor I Receptor (IGF-IR) and Cyclin-dependent kinases (CDKs), and anti-angiogenic compounds having another mechanism of action than decreasing the protein kinase activity.

Compounds which decrease the activity of VEGF are especially compounds which inhibit the VEGF receptor, especially the tyrosine kinase activity of the VEGF receptor, and compounds binding to VEGF, and are in particular those compounds, proteins and monoclonal antibodies generically and specifically disclosed in WO 98/35958 (describing compounds of formula I), WO 00/09495, WO 00/27820, WO 00/59509, WO 98/11223, WO 00/27819, WO 01/55114, WO 01/58899 and EP 0 769 947; those as described by M. Prewett et al in Cancer Research 59 (1999) 5209-5218, by F. Yuan et al in Proc. Natl. Acad. Sci. USA, vol. 93, pp. 14765-14770,

December 1996, by Z. Zhu et al in Cancer Res. 58, 1998, 3209-3214, and by J. Mordenti et al in Toxicologic Pathology, vol. 27, no. 1, pp 14-21, 1999; in WO 00/37502 and WO 94/10202; Angiostatin™, described by M. S. O'Reilly et al, Cell 79, 1994, 315-328; and Endostatin™, described by M. S. O'Reilly et al, Cell 88, 1997, 277-285;

compounds which decrease the activity of EGF are especially compounds which inhibit the EGF receptor, especially the tyrosine kinase activity of the EGF receptor, and compounds binding to EGF, and are in particular those compounds generically and specifically disclosed in WO 97/02266 (describing compounds of formula IV), EP 0 564 409, WO 99/03854, EP 0520722, EP 0 566 226, EP 0 787 722, EP 0 837 063, WO 98/10767, WO 97/30034, WO 97/49688, WO 97/38983 and, especially, WO 96/33980;

compounds which decrease the activity of c-Src include, but are not limited to, compounds inhibiting the c-Src protein tyrosine kinase activity as defined below and to SH2 interaction inhibitors such as those disclosed in WO97/07131 and WO97/08193;

compounds inhibiting the c-Src protein tyrosine kinase activity include, but are not limited to, compounds belonging to the structure classes of pyrrolopyrimidines, especially pyrrolo[2,3-d]pyrimidines, purines, pyrazopyrimidines, especially pyrazo[3,4-d]pyrimidines, pyrazopyrimidines, especially pyrazo[3,4-d]pyrimidines and pyridopyrimidines, especially pyrido[2,3-d]pyrimidines. Preferably, the term relates to those compounds disclosed in WO 96/10028, WO 97/28161, WO97/32879 and WO97/49706;

compounds which decrease the activity of the protein kinase C are especially those staurosporine derivatives disclosed in EP 0 296 110 (pharmaceutical preparation described in WO 00/48571) which compounds are protein kinase C inhibitors;

further specific compounds that decrease protein kinase activity and which may also be used in combination with the compounds of the present invention are Imatinib (Gleevec®/Glivec®), PKC412, Iressa™ (ZD1839), PKI166, PTK787, ZD6474, GW2016, CHIR-200131, CEP-7055/CEP-5214, CP-547632 and KRN-633;

anti-angiogenic compounds having another mechanism of action than decreasing the protein kinase activity include, but are not limited to e.g. thalidomide (THALOMID), celecoxib (Celebrex), SU5416 and ZD6126.

The term "gonadorelin agonist" as used herein includes, but is not limited to abarelix, goserelin and goserelin acetate. Goserelin is disclosed in US 4,100,274.

The term "anti-androgens" as used herein includes, but is not limited to bicalutamide (CASODEX™), which can be formulated, e.g. as disclosed in US 4,636,505.

The term "bengamides" relates to bengamides and derivatives thereof having aniproliferative properties.

The term "bisphosphonates" as used herein includes, but is not limited to etridonic acid, clodronic acid, tiludronic acid, pamidronic acid, alendronic acid, ibandronic acid, risedronic acid and zoledronic acid.

The term "antiproliferative antibodies" as used herein includes, but is not limited to trastuzumab (Herceptin™), Trastuzumab-DM1, erlotinib (Tarceva™), bevacizumab (Avastin™), rituximab (Rituxan®), PRO64553 (anti-CD40) and 2C4 Antibody.

The structure of the active agents identified by code nos., generic or trade names may be taken from the actual edition of the standard compendium "The Merck Index" or from databases, e.g. Patents International (e.g. IMS World Publications).

In accordance with the foregoing the present invention provides in a yet further aspect:

(6) A method as defined above comprising co-administration, e.g. concomitantly or in sequence, of a therapeutically effective amount of a) a compound of formula I or a pharmaceutically acceptable salt thereof, and b) a second drug substance, said second drug substance being for example for use in any of the particular indications hereinbefore set forth.

(7) A combination comprising a therapeutically effective amount of a ZAP-70, FAK and/or Syk tyrosine kinase inhibitor, e.g. a compound of formula I or a pharmaceutically acceptable salt thereof, and a second drug substance, said second drug substance being for example as disclosed above.


Where a ZAP-70, FAK and/or Syk tyrosine kinase inhibitor, e.g. a compound of formula I, is administered in conjunction with other immunosuppressive/immunomodulatory, anti-inflammatory or antineoplastic agent, e.g. as disclosed above, dosages of the co-administered drug or agent will

of course vary depending on the type of co-drug or –agent employed, or the specific drug or agent used, or the condition being treated and so forth.

Representative FAK inhibitors are the compounds of Examples Nos. 187-203 and 209-212.

Claims:

1. A compound of formula I



wherein

X is =CR⁰- or =N-;

each of R⁰, R¹, R², R³ and R⁴ independently is hydrogen; hydroxy; C₁-C₈alkyl; C₂-C₈alkenyl;

C₃-C₈cycloalkyl; C₃-C₈cycloalkyl-C₁-C₈alkyl; hydroxyC₁-C₈alkyl; C₁-C₈alkoxyC₁-C₈alkyl; hydroxyC₁-C₈alkoxyC₁-C₈alkyl; arylC₁-C₈alkyl which optionally may be substituted on the ring by hydroxy, C₁-C₈alkoxy, carboxy or C₁-C₈alkoxycarbonyl;

or R³ and R⁴ form together with the nitrogen and carbon atoms to which they are attached a 5 to 10 membered heterocyclic ring and comprising additionally 1, 2 or 3 heteroatoms selected from N, O and S;

or each of R¹, R² and R³, independently, is halogen; halo-C₁-C₈alkyl; C₁-C₈alkoxy; halo-C₁-C₈alkoxy; hydroxyC₁-C₈alkoxy; C₁-C₈alkoxyC₁-C₈alkoxy; aryl; arylC₁-C₈alkoxy; heteroaryl; heteroaryl-C₁-C₄alkyl; 5 to 10 membered heterocyclic ring; nitro; carboxy; C₂-C₈alkoxycarbonyl; C₂-C₈alkylcarbonyl; -N(C₁-C₈alkyl)C(O)C₁-C₈alkyl; -N(R¹⁰)R¹¹; -CON(R¹⁰)R¹¹; -SO₂N(R¹⁰)R¹¹; or -C₁-C₄-alkylene-SO₂N(R¹⁰)R¹¹; wherein each of R¹⁰ and R¹¹ independently is hydrogen; hydroxy; C₁-C₈alkyl; C₂-C₈alkenyl; C₃-C₈cycloalkyl; C₃-C₈cycloalkyl-C₁-C₈alkyl; C₁-C₈alkoxyC₁-C₈alkyl; hydroxyC₁-C₈alkoxyC₁-C₈alkyl; hydroxyC₁-C₈alkyl; (C₁-C₈alkyl)-carbonyl; arylC₁-C₈alkyl which optionally may be substituted on the ring by hydroxy, C₁-C₈alkoxy, carboxy or C₂-C₈alkoxycarbonyl; or 5 to 10 membered heterocyclic ring;

or R¹ and R² form together with the C-atoms to which they are attached aryl or a 5 to 10 membered heteroaryl residue comprising one or two heteroatoms selected from N, O and S; or

each of R⁵ and R⁶ independently is hydrogen; halogen; cyano; C₁-C₈alkyl; halo-C₁-C₈alkyl; C₂-C₈alkenyl; C₂-C₈alkynyl; C₃-C₈cycloalkyl; C₃-C₈cycloalkylC₁-C₈alkyl; C₅-C₁₀arylC₁-C₈alkyl;


each of R⁷, R⁸ and R⁹ is independently hydrogen; hydroxy; C₁-C₈alkyl; C₂-C₈alkenyl; halo-C₁-C₈alkyl; C₁-C₈alkoxy; C₃-C₈cycloalkyl; C₃-C₈cycloalkylC₁-C₈alkyl; arylC₁-C₈alkyl;

-Y-R¹² wherein Y is a direct bond or O and R¹² is a substituted or unsubstituted 5, 6 or 7 membered heterocyclic ring comprising 1, 2 or 3 heteroatoms selected from N, O and S; carboxy; (C₁-C₈alkoxy)-carbonyl; -N(C₁₋₈alkyl)-CO-NR¹⁰R¹¹; -CONR¹⁰R¹¹; -N(R¹⁰)(R¹¹); -SO₂N(R¹⁰)R¹¹; R⁷ and R⁸ or R⁸ and R⁹, respectively form together with the carbon atoms to which they are attached, a 5 or 6 membered heteroaryl comprising 1, 2 or 3 heteroatoms selected from N, O and S; or a 5 or 6 membered carbocyclic ring.

in free form or salt form.

2. A compound according to claim 1 wherein at most one of R¹, R² or R³ is -CON(R¹⁰)R¹¹; or -SO₂N(R¹⁰)R¹¹.

3. A process for the production of a compound of formula I according to claim 1, comprising the steps of reacting a compound of formula II



wherein R¹, R², R³, R⁴, R⁵, R⁶ and X are as defined in claim 1, and Y is a leaving group;

with a compound of formula III



wherein R⁷, R⁸ and R⁹ are as defined in claim 1;

and recovering the resulting compound of formula I in free form or in salt form, and, where required, converting the compound of formula I obtained in free form into the desired salt form, or vice versa.

4. A compound according to claim 1 in free form or in pharmaceutically acceptable salt form, for use as a pharmaceutical.
5. A pharmaceutical composition comprising a compound of formula I according to claim 1 or a pharmaceutically acceptable salt thereof, together with one or more pharmaceutically acceptable carriers or diluents therefor.
6. The use of a compound of formula I according to claim 1 in free form or in pharmaceutically acceptable salt form, as a pharmaceutical for the treatment or prevention of a disease or condition in which ZAP-70, FAK and/or Syk tyrosine kinase activation plays a role or is implicated.
7. The use of a compound of formula I according to claim 1 in free form or in pharmaceutically acceptable salt form, as a pharmaceutical for the treatment or prevention of a disease or condition in which ZAP-70, FAK and/or Syk tyrosine kinase activation plays a role or is implicated.
8. A combination which comprises (a) a therapeutically effective amount of a ZAP-70, FAK and/or Syk inhibitor and (b) a second drug substance.
9. A method for treating or preventing a disease or condition in which ZAP-70, FAK and/or Syk tyrosine kinase activation plays a role or is implicated, in a subject in need of such treatment, which comprises administering to such subject a therapeutically effective amount of a compound of formula I according to claim 1 or a pharmaceutically acceptable salt thereof.
10. A method for treating or preventing a disease or condition in which ZAP-70, FAK and/or Syk tyrosine kinase activation plays a role or is implicated, in a subject in need of such treatment, which comprises administering to such subject a therapeutically effective amount of a ZAP-70, FAK and/or Syk inhibitor in combination with a second drug substance.

SEQUENCE LISTING

<110> Novartis AG

<120> Pyrimidine derivatives

<130> 4-32366A

<150> GB 0206215.6

<151> 2002-03-15

<160> 4

<170> PatentIn version 3.1

<210> 1

<211> 15

<212> PRT

<213> Artificial sequence

<220>

<223> LAT-11 is an artificial sequence for use in the ZAP-70 kinase assay

<220>

<221> MISC_FEATURE

<222> (1)..(1)

<223> X=L(+)-biotinyl-aminohexanoyl

<400> 1

Xaa Glu Glu Gly Ala Pro Asp Tyr Glu Asn Leu Gln Gln Leu Asn
1 5 10 15

<210> 2

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Biot-Y397 is a sequence for use in FAK assay

<220>

<221> MISC_FEATURE

<222> (1)..(1)

<223> Biotin

<400> 2

Xaa Ser Glu Thr Asp Asp Tyr Ala Glu Ile Ile Asp
1 5 10

<210> 3

<211> 21

<212> DNA

<213> Artificial sequence

<220>

<223> PCR primer for preparing human FAK cDNA

<400> 3
atggcagctg cttaccttga c

21

<210> 4

<211> 21

<212> DNA

<213> Artificial sequence

<220>

<223> PCR primer for preparing human FAK cDNA

<400> 4
tcagtgtggt ctcgtctgcc c

21

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/02710

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D239/48 A61K31/505 A61P35/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 6 235 746 B1 (DAVIS JEREMY MARTIN ET AL) 22 May 2001 (2001-05-22) claims; examples 89-127 ----	1,3-10
X	WO 97 19065 A (CELLTECH THERAPEUTICS LTD ;DAVIS PETER DAVID (GB); MOFFAT DAVID FE) 29 May 1997 (1997-05-29) claims; examples 89-127 ----	1,3-10
X	WO 01 60816 A (AMGEN INC) 23 August 2001 (2001-08-23) claims; examples 56-60 ----	1,3-10
X	EP 1 184 376 A (YAMANOUCHI PHARMA CO LTD) 6 March 2002 (2002-03-06) claims; examples 73-108 ----	1-10
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

11 June 2003

Date of mailing of the international search report

18/06/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Menegaki, F

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/02710

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 01 64655 A (BREault GLORIA ANNE ;PEASE ELIZABETH JANET (GB); ASTRAZENECA UK LT) 7 September 2001 (2001-09-07) page 50 -page 52; claims; examples 5-13,26,27,30,31 ----	1,3-10
X	WO 00 39101 A (BREault GLORIA ANNE ;PEASE JANET ELIZABETH (GB); ASTRAZENECA UK LT) 6 July 2000 (2000-07-06) page 103 -page 104; claims; examples ----	1,3-10
X	WO 00 12485 A (BREault GLORIA ANNE ;PEASE JANET ELIZABETH (GB); ZENECA LTD (GB)) 9 March 2000 (2000-03-09) page 83 -page 84; claims; examples 83-1,83-2 ----	1,3-10
X	EP 1 054 004 A (YAMANOUCHI PHARMA CO LTD) 22 November 2000 (2000-11-22) claims; examples 7,8,48; tables 3,5 ----	1,3-10
P,X	WO 03 018021 A (AMGEN INC) 6 March 2003 (2003-03-06) claims; examples -----	1,3-10

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 03/02710

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 6,7,9 and 10 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compounds/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/02710

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 6235746	B1	22-05-2001	AU 7631496 A DE 69627179 D1 EP 0862560 A1 WO 9719065 A1 US 5958935 A	11-06-1997 08-05-2003 09-09-1998 29-05-1997 28-09-1999
WO 9719065	A	29-05-1997	AU 7631496 A DE 69627179 D1 EP 0862560 A1 WO 9719065 A1 US 6235746 B1 US 5958935 A	11-06-1997 08-05-2003 09-09-1998 29-05-1997 22-05-2001 28-09-1999
WO 0160816	A	23-08-2001	AU 3704101 A CA 2400447 A1 EP 1257546 A1 WO 0160816 A1 US 2002052386 A1	27-08-2001 23-08-2001 20-11-2002 23-08-2001 02-05-2002
EP 1184376	A	06-03-2002	AU 5107900 A EP 1184376 A1 WO 0075113 A1 JP 2001055378 A	28-12-2000 06-03-2002 14-12-2000 27-02-2001
WO 0164655	A	07-09-2001	AU 3397601 A BR 0108834 A CA 2398685 A1 CN 1406230 T EP 1268444 A1 WO 0164655 A1 NO 20024153 A	12-09-2001 10-12-2002 07-09-2001 26-03-2003 02-01-2003 07-09-2001 29-10-2002
WO 0039101	A	06-07-2000	AU 1874300 A BR 9916590 A CA 2352896 A1 CN 1335838 T EP 1140860 A1 WO 0039101 A1 JP 2002533446 T NO 20013038 A	31-07-2000 23-10-2001 06-07-2000 13-02-2002 10-10-2001 06-07-2000 08-10-2002 22-08-2001
WO 0012485	A	09-03-2000	AU 5438299 A EP 1107957 A1 WO 0012485 A1 JP 2002523497 T	21-03-2000 20-06-2001 09-03-2000 30-07-2002
EP 1054004	A	22-11-2000	AU 1507199 A EP 1054004 A1 US 6432963 B1 WO 9931073 A1	05-07-1999 22-11-2000 13-08-2002 24-06-1999
WO 03018021	A	06-03-2003	WO 03018021 A1	06-03-2003